


## CLAIMS:

What is claimed is:

1. A method of treating chronic or neuropathic pain, treating or preventing  
 5 migraine headache, or treating urge, stress or mixed urinary incontinence comprising  
 administration of an effective amount of a compound of formula IA-IF having the  
 following structure:



IA-IF

wherein:

10

the carbon atom designated \* is in the R or S configuration;

R<sup>1</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or C<sub>4</sub>-C<sub>7</sub>  
 cycloalkylalkyl, each of which is optionally substituted with 1 to 3 substituents

15 independently selected at each occurrence thereof from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, aryl, -  
 CN, -OR<sup>9</sup> and -NR<sup>9</sup>R<sup>10</sup>;

R<sup>2</sup> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>4</sub>-C<sub>7</sub>  
 cycloalkylalkyl or C<sub>1</sub>-C<sub>6</sub> haloalkyl;

20

R<sup>3</sup> is H, halogen, -OR<sup>11</sup>, -S(O)R<sup>12</sup>, -S(O)<sub>n</sub>NR<sup>11</sup>R<sup>12</sup>, -CN, -C(O)R<sup>12</sup>, -C(O)NR<sup>11</sup>R<sup>12</sup>,  
 C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl, -  
 O(phenyl) or -O(benzyl), wherein each of -O(phenyl) and -O(benzyl) is optionally  
 substituted from 1 to 3 times with a substituent selected independently at each

- occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy, or wherein R<sup>3</sup> is a C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl group, then said group is optionally substituted with from 1 to 3 substituents selected independently at each occurrence thereof from C<sub>1</sub>-C<sub>3</sub> alkyl,
- 5 halogen, aryl, -CN, -OR<sup>9</sup> and -NR<sup>9</sup>R<sup>10</sup>;
- provided that for compounds of formula IA, R<sup>3</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl, each of which is optionally substituted with from 1 to 3 substituents selected independently at each occurrence thereof from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, aryl, -CN, -OR<sup>9</sup> and -NR<sup>9</sup>R<sup>10</sup>;
- 10 provided that for compounds of formula IB, R<sup>3</sup> is - O(phenyl), -O(benzyl), -OC(O)R<sup>13</sup> or -S(O)<sub>n</sub>R<sup>12</sup>, each of -O(phenyl) and -O(benzyl) is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy;
- 15 R<sup>4</sup> is H, halogen, -OR<sup>11</sup>, -S(O)<sub>n</sub>R<sup>12</sup>, -S(O)NR<sup>11</sup>R<sup>12</sup>, -CN, -C(O)R<sup>12</sup>, -C(O)NR<sup>11</sup>R<sup>12</sup>, -NR<sup>11</sup>R<sup>12</sup>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl, O(phenyl) or -O(benzyl), wherein each of -O(phenyl) and -O(benzyl) is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl or C<sub>1</sub>-C<sub>4</sub> alkoxy and wherein R<sup>4</sup> is a C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl group, then said group is optionally substituted with from 1 to 3 substituents selected independently at each occurrence thereof from C<sub>1</sub>-C<sub>3</sub> alkyl,
- 20 halogen, aryl, -CN, -OR<sup>9</sup> and -NR<sup>9</sup>R<sup>10</sup>;
- provided that for compounds of formula IC, R<sub>4</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, or C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl, each of which is optionally substituted with from 1 to 3 substituents selected independently at each occurrence thereof from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, aryl, -CN, -OR<sup>9</sup> and -NR<sup>9</sup>R<sup>10</sup>, or R<sup>5</sup> and R<sup>6</sup> or R<sup>6</sup> and R<sup>7</sup> may be -O-C(R<sup>12</sup>)<sub>2</sub>-O-; provided that for compounds of formula ID, R<sup>4</sup> is - O(phenyl), -O(benzyl), -OC(O)R<sup>13</sup>, -NR<sup>11</sup>R<sup>12</sup> or -S(O)<sub>n</sub>R<sup>12</sup>, each of -O(phenyl) and -
- 25 O(benzyl) is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy;
- 30

- $R^5$ ,  $R^6$  and  $R^7$  in compounds of each of the formulae IA, IB, IC, ID, IE and IF are each independently H, halogen,  $-OR^{11}$ ,  $-S(O)_nR^{12}$ ,  $-CN$ ,  $-C(O)R^{12}$ ,  $-NR^{11}R^{12}$ ,  $-C(O)NR^{11}R^{12}$ ,  $-NR^{11}C(O)R^{12}$ ,  $-NR^{11}C(O)_2R^{12}$ ,  $-NR^{11}C(O)NR^{12}R^{13}$ ,  $C_1$ - $C_6$  alkyl,  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_6$  alkynyl,  $C_3$ - $C_6$  cycloalkyl or  $C_4$ - $C_7$  cycloalkylalkyl, wherein each of  $R^5$ ,  $R^6$  and  $R^7$  is a  $C_1$ - $C_6$  alkyl,  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_6$  alkynyl,  $C_3$ - $C_6$  cycloalkyl or  $C_4$ - $C_7$  cycloalkylalkyl group, then said group is optionally substituted with from 1 to 3 substituents selected independently at each occurrence thereof from  $C_1$ - $C_3$  alkyl, halogen, aryl,  $-CN$ ,  $-OR^9$  and  $-NR^9R^{10}$ , or  $R^5$  and  $R^6$  or  $R^6$  and  $R^7$  may be  $-O-C(R^{12})_2-O-$ ;
- provided that for compounds of formula IE at least one of  $R^5$  or  $R^7$  is fluoro, chloro, or methyl;
- or  $R^7$  and  $R^6$  are each independently  $-O-C(R^{12})_2-O-$  in compounds of the formulae IE, but only where  $R^2$  is fluoro, chloro or methyl;
- or  $R^7$  and  $R^6$  can independently also be  $-O-C(R^{12})_2-O-$  in compounds of the formulae IE, but only where  $R^7$  is fluoro, chloro or methyl;
- $R^8$  is H, halogen, or  $OR^{11}$ , provided that for compounds of formula IF,  $R^8$  is halogen,  $R^9$  and  $R^{10}$  are each independently H,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  haloalkyl,  $C_1$ - $C_4$  alkoxyalkyl,  $C_3$ - $C_6$  cycloalkyl,  $C_4$ - $C_7$  cycloalkylalkyl,  $-C(O)R^{13}$ , phenyl or benzyl, where phenyl or benzyl is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  haloalkyl, or  $C_1$ - $C_4$  alkoxy;
- or  $R^9$  and  $R^{10}$  are taken together with the nitrogen to which they are attached to form piperidine, pyrrolidine, piperazine, N-methylpiperazine, morpholine, or thiomorpholine;
- $R^{11}$  is H,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  haloalkyl,  $C_1$ - $C_4$  alkoxyalkyl,  $C_3$ - $C_6$  cycloalkyl,  $C_4$ - $C_7$  cycloalkylalkyl,  $-C(O)R^{13}$ , phenyl or benzyl, where  $R^{11}$  is a  $C_1$ - $C_4$  alkyl, phenyl or benzyl group, then said group is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  haloalkyl, or  $C_1$ - $C_4$  alkoxy;
- $R^{12}$  is H, amino,  $C_1$ - $C_4$  alkyl,  $(C_1$ - $C_4$  alkyl)amino,  $C_1$ - $C_4$  haloalkyl,  $C_1$ - $C_4$  alkoxyalkyl,  $C_3$ - $C_6$  cycloalkyl,  $C_4$ - $C_7$  cycloalkylalkyl, phenyl or benzyl, where phenyl or benzyl is optionally substituted from 1 to 3 times with a substituent selected independently from halogen, cyano,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  haloalkyl and  $C_1$ - $C_6$  alkoxy;

or R<sup>11</sup> and R<sup>12</sup> are taken together with the nitrogen to which they are attached to form piperidine, pyrrolidine, piperazine, N-methylpiperazine, morpholine, or thiomorpholine; provided that only one of R<sup>9</sup> and R<sup>10</sup> or R<sup>9</sup> and R<sup>10</sup> are taken together with the  
 5 nitrogen to which they are attached to form piperidine, pyrrolidine, piperazine, N-methylpiperazine, morpholine, or thiomorpholine;

R<sup>13</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl or phenyl;

n is 0, 1, or 2, and;

10

aryl is phenyl which is optionally substituted 1-3 times with halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl and C<sub>1</sub>-C<sub>4</sub> alkoxy,

or an oxide thereof, a pharmaceutically acceptable salt thereof, a solvate thereof, or  
 15 prodrug thereof.

2. A method of claim 1, wherein R<sup>1</sup> is C<sub>1</sub>-C<sub>3</sub> alkyl.

3. A method of claim 2, wherein R<sup>1</sup> is CH<sub>3</sub>.

20

4. A method of claim 1, wherein R<sup>2</sup> is H, C<sub>1</sub>-C<sub>4</sub> alkyl or C<sub>1</sub>-C<sub>6</sub> haloalkyl.

5. A method of claim 4, wherein R<sup>2</sup> is H or CH<sub>3</sub>.

25 6. A method of claim 1, wherein R<sup>3</sup> is H or R<sup>3</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl, each of which is optionally substituted with from 1 to 3 substituents selected independently at each occurrence thereof from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, aryl, -CN, -OR<sup>9</sup> and NR<sup>9</sup>R<sup>10</sup>, or R<sup>3</sup> is -O(phenyl) or -O(benzyl) optionally substituted from 1 to 3 times with a substituent selected independently at each  
 30 occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy.

7. A method of claim 6, wherein R<sup>3</sup> is methyl, ethyl, propyl, or isopropyl.

8. A method of claim 6, wherein  $R^3$  is -O(phenyl) or -O-CH<sub>2</sub>-(phenyl), each of which is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy.
- 5
9. A method of claim 6, wherein  $R^3$  is H.
10. A method of claim 1, wherein  $R^4$  is H, or  $R^4$  is -NR<sup>11</sup>R<sup>12</sup> or  $R^4$  is C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl, each of which is optionally substituted, or  
10 wherein  $R^4$  is -O(phenyl) or -O(benzyl), each of which is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy.
11. A method of claim 10, wherein  $R^4$  is methyl, ethyl, propyl, or isopropyl.
- 15
12. A method of claim 10, wherein  $R^4$  is -O(phenyl) or -O(CH<sub>2</sub>)phenyl, each of which is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy.
- 20
13. A method of claim 10, wherein  $R^4$  is H.
14. A method of claim 1, wherein  $R^3$  and  $R^4$  are each H or wherein  $R^3$  and  $R^4$  are each halogen.
- 25
15. A method of claim 1, wherein one of  $R^3$  and  $R^4$  is H and the other is CH<sub>3</sub>.
16. A method of claim 1, wherein  $R^5$ ,  $R^6$  and  $R^7$  are each H, halogen, -OR<sup>11</sup>, -NR<sup>11</sup>R<sup>12</sup>, C<sub>1</sub>-C<sub>6</sub> alkyl and substituted C<sub>1</sub>-C<sub>6</sub> alkyl.
- 30
17. A method of claim 16, wherein  $R^5$ ,  $R^6$  and  $R^7$  are each H.

18. A method of claim 16, wherein one of  $R^5$  or  $R^7$  is F, Cl or Me and the other of  $R^5$  or  $R^7$  and  $R^6$  are H, halogen,  $-OR^{11}$ ,  $-NR^{11}R^{12}$ , or optionally substituted  $C_1$ - $C_6$  alkyl.
19. A method of claim 18, wherein  $R^5$  is F, Cl or Me; and  $R^7$  is H.
20. The method of claim 18, wherein  $R^5$  is F, Cl or Me; and  $R^6$  is H.
21. A method of claim 1, wherein  $R^8$  is halogen.
22. A method of claim 21, wherein  $R^8$  is fluoro.
23. A method of claim 1, wherein:
- $R^1$  is  $C_1$ - $C_3$  alkyl;
- $R^2$  is H,  $C_1$ - $C_4$  alkyl or  $C_1$ - $C_6$  haloalkyl;
- $R^3$  is  $C_1$ - $C_4$  alkyl,  $C_3$ - $C_6$  cycloalkyl or  $C_4$ - $C_7$  cycloalkylalkyl, each of which is optionally substituted, or  $R^3$  is  $-O(\text{phenyl})$  or  $-O(\text{benzyl})$ , each of which is optionally substituted, or  $R^3$  is H;  $R^4$  is H,  $C_1$ - $C_4$  alkyl,  $C_3$ - $C_6$  cycloalkyl or  $C_4$ - $C_7$  cycloalkylalkyl, each of which is optionally substituted with from 1 to 3 substituents selected independently at each occurrence thereof from  $C_1$ - $C_3$  alkyl, halogen, aryl,  $-CN$ ,  $-OR^9$  and  $-NR^9R^{10}$ , or  $R^4$  is  $-NR^{11}R^{12}$ ,  $-O(\text{phenyl})$  or  $-O(\text{benzyl})$ , wherein said  $-O(\text{phenyl})$  or  $-O(\text{benzyl})$ , is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  haloalkyl, or  $C_1$ - $C_4$  alkoxy;
- or  $R^3$  and  $R^4$  are each halogen;
- $R^5$ ,  $R^6$  and  $R^7$  are each H, halogen,  $-OR^{11}$ ,  $-NR^{11}R^{12}$ , optionally substituted  $C_1$ - $C_6$  alkyl, or one of  $R^5$  and  $R^7$  is Cl, F or Me and the other of  $R^5$  and  $R^7$  and  $R^6$  is H, halogen,  $-OR^{11}$ ,  $-NR^{11}R^{12}$ ,  $C_1$ - $C_6$  alkyl or substituted  $C_1$ - $C_6$  alkyl.
24. A method of claim 23, wherein:

R<sup>1</sup> is CH<sub>3</sub>;

R<sup>2</sup> is H or CH<sub>3</sub>;

R<sup>3</sup> is H, F, methyl, ethyl, propyl, isopropyl, -O(phenyl) or -O-CH<sub>2</sub>-(phenyl), wherein said - O(phenyl) or -O-CH<sub>2</sub>-(phenyl) is optionally substituted from 1 to 3 times with a  
 5 substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy;

R<sup>4</sup> is H, F methyl, ethyl, propyl, isopropyl, -O(phenyl) or -O-CH<sub>2</sub>-(phenyl), wherein said - O(phenyl) or -O-CH<sub>2</sub>-(phenyl) is optionally substituted from 1 to 3 times with a  
 10 substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy;

R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> are each H or R<sup>5</sup> is F, Cl or Me, or one of R<sup>6</sup> or R<sup>7</sup> is H and the other of R<sup>6</sup> and R<sup>7</sup> is halogen, -OR<sup>11</sup>, -NR<sup>11</sup>R<sup>12</sup>, or optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl.

15

25. A method of claim 23, wherein R<sup>8</sup> is halogen.

26. A method according to claim 1, wherein the carbon atom designated \* is in the R configuration.

20

27. A method according to claim 1, wherein the carbon atom designated \* is in the S configuration.

28. A method comprising a mixture of stereoisomeric compounds of claim 1  
 25 wherein the carbon atom designated \* is in the S or R configuration.

29. A method according to claim 1, wherein the compound is selected from the group:

30 2,7-dimethyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;

4-(4-methoxy)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;

2,7-dimethyl-4-(4-fluoro)phenyl-1,2,3,4-tetrahydroisoquinoline;

2,7-dimethyl-4-(3-fluoro)phenyl-1,2,3,4-tetrahydroisoquinoline;

5 4-(3,4-difluoro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;

2,7-dimethyl-4-(4-fluoro-3-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;

4-(3-chloro-4-fluoro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;

10

4-(3-chloro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;

2,7-dimethyl-4-(4-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;

15 2,7-dimethyl-4-(3-fluoro-4-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;

4-(4-chloro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;

4-(4-chloro-3-fluoro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;

20

4-(3,4-dichloro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;

7-ethyl-2-methyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;

25 4-(3,4-difluoro)phenyl-7-ethyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

7-fluoro-4-(4-methoxy)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

7-fluoro-4-(3-fluoro-4-methoxy)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

30

7-fluoro-4-(3-fluoro-4-methyl)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

7-fluoro-4-(4-chloro-3-fluoro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;



- 4-(3,4-difluoro)phenyl-7-fluoro-2-methyl-1,2,3,4-tetrahydroisoquinoline;
- 4-(3-chloro)phenyl-7-fluoro-2-methyl-1,2,3,4-tetrahydroisoquinoline;
- 5 7-cyano-2-methyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;
- 2-methyl-4-phenyl-7-trifluoromethyl-1,2,3,4-tetrahydroisoquinoline;
- 10 4-phenyl-1,2,7-trimethyl-1,2,3,4-tetrahydroisoquinoline;
- 4-(4-chloro)phenyl-1,2-dimethyl-1,2,3,4-tetrahydroisoquinoline;
- 4-(3,4-difluoro)phenyl-1,2-dimethyl-1,2,3,4-tetrahydroisoquinoline;
- 15 4-phenyl-2,7,8-trifluoromethyl-1,2,3,4-tetrahydroisoquinoline;
- 2,7-dimethyl-8-fluoro-4-phenyl-1,2,3,4-tetrahydroisoquinoline;
- 20 2,8-dimethyl-7-fluoro-4-phenyl-1,2,3,4-tetrahydroisoquinoline;
- 2,7-dimethyl-8-methoxy-4-phenyl-1,2,3,4-tetrahydroisoquinoline;
- 2,7-dimethyl-8-hydroxy-4-phenyl-1,2,3,4-tetrahydroisoquinoline;
- 25 2-methyl-4-phenyl-7-trifluoromethoxy-1,2,3,4-tetrahydroisoquinoline;
- 4-(3,4-difluoro)phenyl-7-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;
- 30 4-(4-fluoro-3-methyl)phenyl-7-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;
- 4-(3-fluoro-4-methyl)phenyl-7-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;

7-methoxy-4-(3-methyl)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

2-methyl-7-phenoxy-4-phenyl-1,2,3,4-tetrahydroisoquinoline;

5 7-(4-methoxy)phenoxy-2-methyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;

7-benzyloxy-2-methyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;

7-hydroxy-2-methyl-4-(3-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;

10

4-(3-fluoro-4-methyl)phenyl-7-hydroxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;

4-(4-fluoro-3-methyl)phenyl-7-hydroxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;

15 4-(3,4-difluoro)phenyl-7-hydroxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;

4-(3-cyano)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

2,8-dimethyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;

20

2,8-dimethyl-4-(4-fluoro)phenyl-1,2,3,4-tetrahydroisoquinoline;

4-(3,4-difluoro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;

25 4-(3,5-difluoro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;

2,8-dimethyl-4-(3-fluoro)phenyl-1,2,3,4-tetrahydroisoquinoline;

2,8-dimethyl-4-(4-fluoro-3-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;

30

4-(3-chloro-4-fluoro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;

4-(3,4-dichloro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;

- 4-(3-chloro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;
- 4-(4-chloro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;
- 5 4-(4-chloro-3-fluoro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;
- 2,8- dimethyl-4-(4-methoxy)phenyl-1,2,3,4-tetrahydroisoquinoline;
- 10 4-(4-cyano)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;
- 2,8-dimethyl-4-(4-trifluoromethyl)phenyl-1,2,3,4-tetrahydroisoquinoline;
- 2,8-dimethyl-4-(4-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;
- 15 2-methyl- 8-(N-methylamino)methyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;
- 8-(hydroxy)methyl-2-methyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;
- 20 2-methyl-4-phenyl-8-sulfonamide-1,2,3,4-tetrahydroisoquinoline;
- 2-methyl-8-(N-methyl)sulfonamide-4-phenyl-1,2,3,4-tetrahydroisoquinoline;
- 8-methoxy-2-methyl-4-(4-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;
- 25 4-(3,5-difluoro)phenyl-8-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;
- 4-(3-chloro)phenyl-8-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;
- 30 4-(3,4-dichloro)phenyl-8-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;
- 4-(4-chloro-3-fluoro)phenyl-8-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;

4-(3-chloro-4-fluoro)phenyl-8-methoxy-2-methyl-1, 2,3,4-tetrahydroisoquinoline;

4-(3,5-difluoro)phenyl-2-methyl-1,2,3,4- tetrahydroisoquinoline;

5 4-(3-chloro-5-fluoro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

4-(3,5-difluoro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;

4-(3-chloro-5-fluoro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;

10

2-methyl-4-(3,4,5-trifluoro)phenyl-1,2,3,4-tetrahydroisoquinoline;

4-(3- fluoro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

15 4-(3-fluoro-4-methyl)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

4-(4-fluoro-3-methyl)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

4-(3,4-difluoro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

20

4-(3-chloro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

4-(4-chloro-3-fluoro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

25 4-(3-chloro-4- fluoro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

4-(3-cyano)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;

4-(4-acetanilide)-2-methyl-1,2,3,4-tetrahydroisoquinoline;

30

4-(4-chloro)phenyl-4-fluoro-2-methyl-1,2,3,4-tetrahydroisoquinoline;

(3,5-difluoro)-4-phenyl-1,2,7-trimethyl-1,2,3,4-tetrahydroisoquinoline;

(8-fluoro-2-methyl-4-phenyl-1,2,3,4-tetrahydro-7-isoquinoliny)-N-methylmethanamine;

- 5 (2-methyl-4-phenyl-7-isoquinoliny)-N-methylmethanamine;

N-methyl-(2-methyl-4-phenyl-7-isoquinoliny)-N-methylmethanamine;

8-hydroxy-2-methyl-4-phenyl-1,2,3,4-tetrahydro-7-isoquinolinecarbonitrile;

10

(2-methyl-4-phenyl-1,2,3,4-tetrahydro-7-isoquinoliny)methanol; and

2-ethyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline; and

- 15 an oxide thereof, a pharmaceutically acceptable salt thereof, a solvate thereof, or prodrug thereof.